

WEEKLY TEST TYM-02 TEST - 8 RAJPUR ROAD
SOLUTION Date 22-09-2019

[PHYSICS]

1. According to free body diagram of block A,

$$F - T_1 = m_1 a \quad \dots (i)$$

$$T_1 - T_2 = m_2 a \quad \dots (ii)$$

$$T_2 = m_3 a \quad \dots (iii)$$

Adding all the three eqns., we get

$$F = (m_1 + m_2 + m_3)a \quad \text{or} \quad a = \frac{F}{m_1 + m_2 + m_3}$$

$$= \frac{14}{4 + 2 + 1}$$

Putting in eqn. (i), contact force between A and B is

$$T_1 = F - m_1 a = 14 - 4 \times 2 = 6 \text{ N}$$

Hence, correct option is (a)

2. Time period of a simple pendulum is given :

$$T = 2\pi\sqrt{\frac{l}{g}} \quad \text{or} \quad T \propto \sqrt{\frac{l}{g}}$$

When the elevator is accelerating downwards, then net gravitational acceleration is $(g - a)$. So, the time period when elevator is accelerating downwards, is greatest.

3. As per Newton's third law of motion, when a horse pulls a wagon, the force that causes the horse to move forward is the force the ground exerts on it.

4.
$$F = \frac{d}{dt}(Mv) = v \frac{dM}{dt} + M \frac{dv}{dt}$$

As v is a constant,
$$F = v \frac{dM}{dt}$$

But
$$\frac{dM}{dt} = M \text{ kg/s}$$

- ∴ To keep the conveyer belt moving at v m/s,
 Force needed = vM newton

5.

6.
$$F - Mg = Ma$$


$$8000 = 2000 a$$

- ∴ Acceleration is 4 ms^{-2} upwards

7. Considering free-body diagrams of the masses, we have

$$T - 3g = 3a \quad \text{and} \quad 5g - T = 5a$$

Solving for T , we have



$$T = (15/4)g$$

∴ F = Force on the pulley

$$= 2T = 2 \times \frac{15}{4} = 7.5 \text{ kg f}$$

8. Change in momentum in one sec, i.e.,
 F = change in momentum per bullet × no. of bullets fired per second
 = $mv \times n = mnv$

$$9. \quad T_2 = \frac{6}{6+6+6} F = \frac{F}{3}$$

10. For a body to be in equilibrium, it should exist both in translational equilibrium.
 For translational equilibrium, $\Sigma F = 0$
 and for rotational equilibrium, $\Sigma \tau = 0$

11. Acceleration of the mass m_3 = common acceleration of the system = $\frac{F}{\text{total mass}} = \frac{F}{m_1 + m_2 + m_3}$

12. One of the weights given a reading and the other prevents the acceleration of the system. Therefore, the reading is not zero but 10 N.

13. Equations of motion are :

$$F - T_1 = 2a \quad \dots (i)$$

$$T_1 - T_2 = 3a \quad \dots (ii)$$

$$T_2 = 5a \quad \dots (iii)$$


Adding all above equations, we get;

$$F = 10a = 10 \times 1 = 10 \text{ N}$$

14. The tension in the string between P and Q accelerates double the mass as compared to that between A and R. Hence, tension between P and Q = 2 × tension between Q and R


15. $T \cos \theta = T_1 = 10 \times g$
 $T \sin \theta = 98$

$$\therefore \tan \theta = \frac{98}{10 \times 9.8} = 1 \quad \text{or} \quad \theta = 45^\circ$$




CHEMISTRY


16.
17.



- 18.




- 19.


**(2-Propyl) cyclobutane**

20.


The compound can be expanded as

IUPAC name is **3, 4, 4-trimethyloctane**.


21.

**3, 3-Dimethyl cyclohexan-1-ol** is the proper IUPAC name as per 1993 rules. The given answer is as per the old IUPAC rules, still prevalent.


22.

**3-Bromo-1-chlorocyclohexene**


23.

**3-Chloro-4-fluoro-3, 4-dimethylhexane**


24.

**Butane-1,2, 4-tricarboxylic acid**


25.

**3,3-Dimethylbutanoyl chloride**


26.

**3-Cyanopropanamide**


27.

**2-Carbamoylethanoic acid**


28.

**2-(Prop-1-yl) propane-1, 3-dioic acid**

29.

**3-(Chloroformylmethyl) pentane-1, 5-dioyl chloride**

30.

**3-Phenylprop-2-enoic acid**

Its trivial name is cinnamic acid.