

**WEEKLY TEST TYM TEST - 25 Balliwala**  
**SOLUTION Date 20 -10-2019**

**[PHYSICS]**

1. It is the speed of light in free space. Hence, dimension is that of speed, i.e.,  $LT^{-1}$ .

2. Boltzmann's constant = energy/temperature

$$= \frac{[ML^2T^{-2}]}{[\theta]} = [ML^2T^{-2}\theta^{-1}]$$

3. When the body returns to origin, displacement is zero.

$$s = ut + \frac{1}{2}at^2$$

$$0 = 60t - \frac{1}{2} \times 10 \times t^2$$

Solving,  $t = 12 \text{ s}$

4.  $v \propto \lambda^x \rho^y g^z$

Putting dimensions,

$$LT^{-1} = L^x(ML^{-3})^y(LT^{-2})^z$$

Solving, we get  $v \propto \sqrt{g\lambda}$ .

5. Squaring the given equation,

$$A^2 + B^2 + 2\vec{A} \cdot \vec{B} = C^2$$

Moreover,  $A^2 + B^2 = C^2$  ( $\because A = 6, B = 8, C = 10$ )

$\therefore \vec{A} \cdot \vec{B} = 0$ . i.e.,  $\vec{A}$  is  $\perp$  to  $\vec{B}$

6. Putting equations for  $T$  and  $R$ , we get

$$g \left( \frac{2u \sin \alpha}{g} \right)^2 = 2 \times \frac{u^2 \sin 2\alpha}{g}$$

or  $\tan \alpha = 1$  or  $\alpha = 45^\circ$

7. Let  $t$  be the duration of uniform acceleration. Then,  $(9 - t)$  is the retardation. As the velocity at the end of uniform acceleration and at the beginning of retardation is same, we have

$$0 + at = 0 - 2a \times (9 - t)$$

Solving,  $t = 6 \text{ s}$

8. The train is moving with horizontal velocity in a straight line, hence vertical ranges will be same.

For a person inside the train, the horizontal range will be zero, because train is an inertial frame. The coin falls back to his hand. For a person outside the train such as  $C$ , the coin has a horizontal velocity and vertical acceleration  $g$ . Hence, it appears to follow a parabolic path. Hence, he observes a horizontal range.

9.

$$R = \frac{u^2 \sin 2\alpha}{g}$$

i.e.,  $R \propto \sin 2\alpha$  (for a given  $u$ )

$$\frac{R_1}{R_2} = \frac{\sin 30^\circ}{\sin 90^\circ} = \frac{1}{2}$$

$$\therefore R_2 = 2R_1 = 4 \text{ km.}$$

10. The acceleration down the plane =  $\frac{g \sin \theta}{1 + (k^2/r^2)}$

This value is maximum when  $k$  is minimum, which happens for a solid sphere ( $k^2 = \frac{2}{5}r^2$ ).

11. Work done = force  $\times$  displacement =  $100 \times \sin 50^\circ \times 1$ .

20% of this work is used to overcome friction. Hence, energy gained = 80% of this work =  $80 \sin 50^\circ$  Joule.

12. The ball moves towards the left due to inertia to a force towards right. This means the bus is taking a right hand turn. The centripetal force is towards the right. Inertial force on the ball (in this case called centrifugal force) acts towards the left.

13. Potential energy of a satellite =  $-\frac{GMm}{R}$

Kinetic energy of a satellite =  $\frac{GMm}{2R}$

$$\therefore \text{Required ratio} = 2 : 1$$

14. According to Kepler's 3rd law,  $T^2 \propto R^3$ .

Now,  $T_1^2 \propto R^3$  (for a satellite close to the earth)

For a geo-synchronous satellite, its distance from the surface of the earth is  $6R$ , and from centre  $7R$ .

So,  $T_2^2 \propto (7R)^3$


$$\therefore T_2 = 7^{3/2} \cdot T_1 = T_1 7\sqrt{7} \text{ or } \frac{T_2}{T_1} = 7\sqrt{7}$$

15. Before the man hits the ground, he is in a state of free fall (ignoring air resistance). A body in a free fall state experiences weightlessness.

16. Here, the disc has only KE of rotation

$$= \frac{1}{2} I\omega^2 = \frac{1}{2} \times \frac{Mr^2}{2} \frac{v^2}{r^2} = \frac{1}{4} Mv^2$$

17. Change in velocity = final velocity - initial velocity. It is clear from figure, that change in velocity =  $\sqrt{20^2 + 20^2} = 20\sqrt{2}$  m/s along south-west.



18. KE of a satellite =  $\frac{1}{2} |PE|$

$$KE = \frac{GMm}{2R} \text{ and } PE = -\frac{GMm}{R}$$


$$TE = KE + PE = -\frac{GMm}{2R} = -KE = -2 \times 10^6 \text{ J}$$

19. At the highest point, the projectile has only the horizontal component =  $u \cos \alpha = \frac{u}{2}$  ( $\because \alpha = 60^\circ$ )

$$\text{Range} = \frac{u^2}{g} \sin 2\alpha = \frac{2u^2 \sin \alpha \cos \alpha}{g} = \frac{\sqrt{3}u^2}{2g}$$

20. The cylinder begins to slide when the line through centre of gravity  $G$  passes beyond base of cylinder that is when

$$\tan \theta \geq \frac{r}{h/2} = \frac{2r}{h}$$



Hence,  $h = 2r$  ( $\because \theta = 45^\circ$ )  
 $r = \frac{h}{2} = 5 \text{ cm}$

### CHEMISTRY

21. (a)  $\text{Na}_2\text{CO}_3 \cdot x\text{H}_2\text{O}$ . Its weight =  $106 + 18x$ .

Weight of anhydrous  $\text{Na}_2\text{CO}_3 = 106$

$$\% \text{ loss in weight} = \frac{18x \times 100}{106 + 18x} = 63$$

$$\therefore x = 10.27 \approx 10$$

22. (c) In law of reciprocal proportions, the two elements combining with the third element, must combine with each other in the same ratio or multiple of that Ratio of S and O when combine with C is 2 : 1. Ratio of S and O is  $\text{SO}_2$ , is 1 : 1

23. (c) Mol in each case

$$7 \text{ g N}_2 = \frac{7}{28} = 0.25; \quad 2 \text{ g H}_2 = \frac{2}{2} = 1.0;$$

$$16 \text{ g NO}_2 = \frac{16}{46} = 0.34; \quad 16 \text{ g O}_2 = \frac{16}{32} = 0.50$$

Thus hydrogen has maximum moles, hence maximum molecules.

- 24.

$$(a) \lambda = \frac{h}{mv} = \frac{h}{\sqrt{2mE}}$$


$$= \frac{6.6 \times 10^{-34}}{\sqrt{2 \times 1 \times 0.5}} = 6.6 \times 10^{-34}$$




25. (b) 1.  $\left. \begin{array}{l} \text{BO}_3^{3-} \longrightarrow 5 + 8 \times 3 + 3 = 32 \\ \text{CO}_3^{2-} \longrightarrow 6 + 8 \times 3 + 2 = 32 \\ \text{NO}_3^- \longrightarrow 7 + 8 \times 3 + 1 = 32 \end{array} \right\} \text{ISO electronic}$
2.  $\left. \begin{array}{l} \text{SO}_3^{2-} \longrightarrow 16 + 8 \times 3 + 2 = 42 \\ \text{CO}_3^{2-} \longrightarrow 32 \\ \text{NO}_3^- \longrightarrow 32 \end{array} \right\} \text{not ISO electronic}$
3.  $\left. \begin{array}{l} \text{CN}^- \longrightarrow 6 + 7 + 1 = 14 \\ \text{N}_2 \longrightarrow 7 \times 2 = 14 \\ \text{C}_2^{2-} \longrightarrow 6 \times 2 + 2 = 14 \end{array} \right\} \text{ISO electronic}$
4.  $\left. \begin{array}{l} \text{PO}_4^{3-} \longrightarrow 15 \times 8 + 4 + 3 = 50 \\ \text{SO}_4^{2-} \longrightarrow 16 + 8 \times 2 = 50 \\ \text{ClO}_4^- \longrightarrow 17 + 8 \times 4 + 1 = 50 \end{array} \right\} \text{ISO electronic}$

26. (c)  $ns^2 p^1$  is the electronic configuration of III period.  
 $\text{Al}_2\text{O}_3$  is amphoteric oxide

27. (c)  $\text{PF}_5$  trigonal bipyramidal



$\text{BrF}_3$  square pyramidal (distorted)





28. (d)  $\begin{array}{c} (-) \\ \text{:}\ddot{\text{O}}\text{:} - \text{N} \begin{array}{l} \nearrow \ddot{\text{O}}\text{:} \\ \searrow \ddot{\text{O}}\text{:} \end{array} \end{array}$  It has 4 bond pairs and none lone pair on N.

29. (d) The value of  $a$  is a measure of the magnitude of the attractive forces between the molecules of the gas. Greater the value of ' $a$ ', larger is the attractive intermolecular force between the gas molecules. The value of  $b$  related to the effective size of the gas molecules. It is also termed as excluded volume. The gases with higher value of  $a$  and lower value of  $b$  are more liquefiable, hence for  $\text{Cl}_2$  " $a$ " should be greater than for  $\text{C}_2\text{H}_6$ , but for it  $b$  should be less than for  $\text{C}_2\text{H}_6$ .

30. (d)  $\text{NH}_3$  and  $\text{HCl}$  react to form  $\text{NH}_4\text{Cl}$
31. (b) This is combustion reaction, which is always exothermic hence  
 $\Delta H = -ve$   
 As the no. of gaseous molecules are increasing hence entropy increases  
 Now  $\Delta G = \Delta H - T\Delta S$   
 For a spontaneous reaction  
 $\Delta G = -ve$   
 Which is possible in this case as  $\Delta H = -ve$  and  $\Delta S = +ve$ .

32. (c)  $\sigma 1s^2, \sigma^* 1s^2, \sigma 2s^2, \sigma^* 2s^2,$   
 $\sigma 2p_z^2, \pi 2p_x^2, \pi 2p_y^2, \pi^* 2p_x^2, \pi^* 2p_y^2$   
 $\therefore$  No. of antibonding electron pairs = 4

33. (d) The shape of  $\text{BF}_3$  is trigonal planar  and  
 $\mu = 0$  hence it is non polar. The shape of  $\text{NF}_3$  is  
  
 pyramidal  $\delta^-$   $\delta^-$   $\delta^-$  and  $\mu \neq 0$  hence it is polar.

34. 10% (w/w) solution means  $100 \text{ g} = \frac{100}{1.1} \text{ ml}$   
 solution contains 10 g solute,  $\text{NaOH}$   

$$\text{Molarity} = \frac{w \times 1000}{m^l \times v} = \frac{10 \times 1000}{40 \times \left(\frac{100}{1.1}\right)} = 2.75 \text{ M}$$

35. B

36. [a] The average energy per bond in  $\text{O}_2$  is greater than that in  $\text{O}_3$  because dissociation of  $\text{O}_2$  is endothermic