

WEEKLY TEST TYM -01 TEST - 11
SOLUTION Date 30-06-2019

[PHYSICS]

1. Average speed = $\frac{\text{total distance covered}}{\text{total time taken}}$

$$v_{\text{av.}} = \frac{\frac{x}{2} + \frac{x}{2}}{\frac{x/2}{40} + \frac{x/2}{60}} = \frac{x}{\left(\frac{x}{80} + \frac{x}{120}\right)}$$

$$= \frac{80 \times 120}{(120 + 80)} = 48 \text{ km/h}$$

2. $200 = u \times 2 - (1/2) a(2)^2$ or $u - a = 100$ (i)
 $200 + 220 = u(2 + 4) - (1/2)(2 + 4)^2 a$

or $u - 3a = 70$ (ii)

Solving eqns. (i) and (ii), we get; $a = 15 \text{ cm/s}^2$ and $u = 115 \text{ cm/s}$.
 Further, $v = u - at = 115 - 15 \times 7 = 10 \text{ cm/sec.}$

3. When a body slides on an inclined plane, component of weight along the plane produces an acceleration

$$a = \frac{mg \sin \theta}{m} = g \sin \theta = \text{constt.}$$

If s be the length of the inclined plane, then

$$s = 0 + \frac{1}{2} a t^2 = \frac{1}{2} g \sin \theta \times t^2$$

$$\therefore \frac{s'}{s} = \frac{t'^2}{t^2} \text{ or } \frac{s}{s'} = \frac{t^2}{t'^2}$$

$$\text{Given } t = 4 \text{ sec and } s' = \frac{s}{4}$$

$$\therefore t' = t \sqrt{\frac{s'}{s}} = 4 \sqrt{\frac{s}{4s}} = \frac{4}{2} = 2 \text{ sec}$$

4. Given that; $a = 3t + 4$ or $\frac{dv}{dt} = 3t + 4$

$$\therefore \int_0^v dv = \int_0^t (3t + 4) dt \text{ or } v = \frac{3}{2} t^2 + 4t$$

$$v = \frac{3}{2}(2)^2 + 4(2) = 14 \text{ ms}^{-1}$$



5. **For first body :**

$$\frac{1}{2}gt^2 = 176.4 \text{ or } t = \sqrt{\frac{176.4 \times 2}{10}}$$

or $t = 5.9 \text{ s}$

For second body : $t = 3.9 \text{ s}$

$$u(3.9) + \frac{1}{2}g(3.9)^2 = 176.4$$

$$3.9u + \frac{1}{2}(3.9)^2 = 176.4$$

or $u = 24.5 \text{ m/s}$

6. The resultant velocity of the boat and river is $1.0 \text{ km}/0.25 \text{ h} = 4 \text{ km/h}$.

$$\text{Velocity of the river} = \sqrt{5^2 - 4^2} = 3 \text{ km/h}$$

7. Let h be the height of the tower.

Using $v^2 - u^2 = 2as$, we get;

Here, $u = u$, $a = -g$, $s = -h$ and $v = -3u$ (upward direction + ve)

$$\therefore 9u^2 - u^2 = 2gh \text{ or } h = 4u^2/g$$

$$8. t = \sqrt{\frac{2h}{g}}$$

$$s = 10 \times \frac{t}{2} - \frac{1}{2}g \times \frac{t^2}{4} = 5\sqrt{\frac{2h}{g}} - \frac{g}{8} \frac{2h}{g}$$

$$v^2 - u^2 = 2gh \text{ or } 100 = 2gh \text{ or } 10 = \sqrt{2gh}$$

$$s = \sqrt{\frac{2gh \times 2h}{4 \times g}} - \frac{h}{4} = h - \frac{h}{4} = \frac{3h}{4}$$

$$9. t = \frac{1}{u+v} = \frac{1}{\frac{1}{t_1} + \frac{1}{t_2}}$$

$$\text{or } \frac{1}{t} + \frac{1}{t_1} + \frac{1}{t_2} \text{ or } t = \frac{t_1 t_2}{(t_1 + t_2)}$$

10. **For first body :**

$$v^2 = u^2 + 2gh \text{ or } (3)^2 = 0 + 2 \times 9.8 \times h$$

$$\text{or } h = \frac{(3)^2}{2 \times 9.8} = 0.46 \text{ m}$$

For second body :

$$v^2 = (4)^2 + 2 \times 9.8 \times 0.46$$

$$\therefore v = \sqrt{(4)^2 + (2 \times 9.8 \times 0.46)} = 5 \text{ m/s}$$

11. Given $y = 0$

Distance travelled in 10 s,

$$S_1 = \frac{1}{2}a \times 10^2 = 50a$$

Distance travelled in 20 s,

$$S_2 = \frac{1}{2}a \times 20^2 = 200a$$

$$\therefore S_2 = 4S_1$$



12. During the first 5 seconds of the motion, the acceleration is – ve and during the next 5 seconds it becomes positive. (Example : a stone thrown upwards, coming to momentary rest at the highest point). The distance covered remains same during the two intervals of time.
13. Gain in angular KE = loss in PE

$$\text{If } l = \text{length of the pole, moment of inertial of the pole about the edge} = M \left[\frac{l^2}{12} + \frac{l^2}{4} \right] = \frac{Ml^2}{3}$$

$$\text{Loss in potential energy} = \frac{Mgl}{2}$$

$$\text{Gain in angular KE} = \frac{1}{2} I \omega^2 = \frac{1}{2} \times \frac{Ml^2}{3} \times \omega^2$$

$$\therefore \frac{1}{2} \frac{Ml}{3} \omega^2 = \frac{Mgl}{2} \quad \text{or} \quad (I\omega)^2 = 3gl$$

$$\text{or} \quad l\omega = v = \sqrt{3gl}$$

$$= \sqrt{3 \times 10 \times 30} = 30 \text{ ms}^{-1}$$

14. Let the velocity of the scooter be $v \text{ ms}^{-1}$. Then $(v - 10)100 = 100$ or $v = 20 \text{ ms}^{-1}$
15. Let x be the distance between the particles after t second. Then

$$x = vt - \frac{1}{2}at^2 \quad \dots\dots(i)$$

For x to be maximum,

$$\frac{dx}{dt} = 0$$

$$\text{or} \quad v - at = 0$$

$$\text{or} \quad t = \frac{v}{a}$$

Putting this value in eqn. (i), we get;

$$x = v \left(\frac{v}{a} \right) - \frac{1}{2} a \left(\frac{v}{a} \right)^2 = \frac{v^2}{2a}$$

[CHEMISTRY]

- 16.
17. Within a group IE₁ decreases from top to bottom.
18. After the removal of second electron, the ion acquires noble gas configuration and it becomes difficult to remove the third electron.
- 19.
- 20.
- 21.
- 22.
23. With negative sign the chlorine has highest electron gain enthalpy (in magnitude)
- 24.
25. All have 18 electrons.
26. The ionic radii follow the order : C⁴⁻ > N³⁻ > O²⁻ and therefore, N³⁻ would have value between 2.60 and 1.40 Å.
- 27.
- 28.
- 29.
30. The general electronic configuration of d-block elements is $(n - 1)d^{1-10}, ns^{1-2}$. They show variable oxidation state because d-electrons also take part in bond formation. They have take part in bond formation. They have degenerated orbitals. s and p-block elements in general do not show variable oxidation states.

