



**WEEKLY TEST TYM TEST - 33 BALLIWALA**  
**SOLUTION Date 29-12-2019**

**[PHYSICS]**

1. (a)  $v = n\lambda = 2 \times 5 = 10 \text{ cm/sec}$
2. (a)  $v_{\max} = a\omega = a \times 2\pi n = 0.1 \times 2\pi \times 300 = 60\pi \text{ cm/sec}$
3. (c) Phase difference  $= \frac{2\pi}{\lambda} \times \text{path difference}$   
 $\Rightarrow 1.6\pi = \frac{2\pi}{\lambda} \times 40 \Rightarrow \lambda = 50 \text{ cm} = 0.5 \text{ m}$   
 $\Rightarrow v = n\lambda \Rightarrow 330 = 0.5 \times n \Rightarrow n = 660 \text{ Hz}$
4. (a)
5. (a)  $\lambda = \frac{v}{n} = \frac{1.7 \times 1000}{4.2 \times 10^6} = 4 \times 10^{-4} \text{ m}$
6. (c) Velocity of sound in gas  $v = \sqrt{\frac{\gamma RT}{M}} \Rightarrow v \propto \sqrt{\frac{\gamma T}{M}}$   
 $\Rightarrow \frac{v_{N_2}}{v_{He}} = \sqrt{\frac{\gamma_{N_2} \times \frac{M_{He}}{M_{N_2}}}{\gamma_{He} \times \frac{M_{He}}{M_{H_2}}}} = \sqrt{\frac{\frac{7}{5} R \times 4}{\frac{5}{3} R \times 28}} = \frac{\sqrt{3}}{5}$
7. (a) Time required for a point to move from maximum displacement to zero displacement is  $t = \frac{T}{4} = \frac{1}{4n}$   
 $\Rightarrow n = \frac{1}{4t} = \frac{1}{4 \times 0.170} = 1.47 \text{ Hz}$
8. (c)  $\lambda = \frac{v}{n} = \frac{340}{200} = 1.7 \text{ m}$
9. (a) The time taken by the stone to reach the lake  
 $t_1 = \sqrt{\frac{2h}{g}} = \sqrt{\frac{2 \times 500}{10}} = 10 \text{ sec}$  (Using  $h = ut + \frac{1}{2}gt^2$ )  
 Now time taken by sound from lake to the man  
 $t_2 = \frac{h}{v} = \frac{500}{340} \approx 1.5 \text{ sec}$   
 $\Rightarrow \text{Total time} = t_1 + t_2 = 10 + 1.5 = 11.5 \text{ sec.}$
10. (b) Distance between a compression and the nearest rarefaction is  $\frac{\lambda}{2} = 1 \text{ m}$ . Hence  
 $n = \frac{v}{\lambda} = \frac{360}{2} = 180 \text{ Hz}$ .

11. (a)  $v = \sqrt{\frac{\gamma P}{\rho}} \Rightarrow \frac{v_{O_2}}{v_{H_2}} = \sqrt{\frac{\rho_{H_2}}{\rho_{O_2}}} = \sqrt{\frac{1}{16}} = \frac{1}{4}$
12. (d) Speed of sound in gases is  $v = \sqrt{\frac{\gamma RT}{M}} \Rightarrow T \propto M$   
 (Because  $v, \gamma$ -constant). Hence  $\frac{T_{H_2}}{T_{O_2}} = \frac{M_{H_2}}{M_{O_2}}$   
 $\Rightarrow \frac{T_{H_2}}{(273+100)} = \frac{2}{32} \Rightarrow T_{H_2} = 23.2 \text{ K} = -249.7^\circ \text{C}$
13. (c) Path difference  $\Delta = \frac{\lambda}{2\pi} \times \phi \Rightarrow 1 = \frac{\lambda}{2\pi} \times \frac{\pi}{2} \Rightarrow \lambda = 4 \text{ m}$   
 Hence  $v = n\lambda = 120 \times 4 = 480 \text{ m/s}$
14. (a) Suppose the distance between shooter and reflecting surface is  $d$ . Hence time interval for hearing echo is  
  
 $t = \frac{2d}{v} \Rightarrow 8 = \frac{2d}{350} \Rightarrow d = 1400 \text{ m}$
15. (d)  $v = \sqrt{\frac{\gamma P}{\rho}}$ ; as  $P$  changes,  $\rho$  also changes. Hence  $\frac{P}{\rho}$  remains constant so speed remains constant.

16. (b)   
 $2d_1 + 2d_2 = v \times t_1 + v \times t_2 \Rightarrow 2(d_1 + d_2) = v(t_1 + t_2)$   
 $d_1 + d_2 = \frac{v(t_1 + t_2)}{2} = \frac{340 \times (1.5 + 3.5)}{2} = 850 \text{ m}$

17. (b) By using  $v = \sqrt{\frac{\gamma RT}{M}} \Rightarrow v \propto \sqrt{T}$   
 $\frac{v_2}{v_1} = \sqrt{\frac{T_2}{T_1}} = \sqrt{\frac{T+600}{T}} = \sqrt{3} \Rightarrow T = 300 \text{ K} = 27^\circ \text{C}$
18. (c)  $v = \sqrt{\frac{\gamma RT}{M}} \Rightarrow v \propto \sqrt{T}$   
 i.e. if  $v$  is doubled then  $T$  becomes four times,

hence  $T_2 = 4T_1 = 4(273 + 27) = 1200K = 927^\circ C$

19. (c) Since solid has both the properties (rigidity and elasticity)

20. (b) Frequency of wave is  $n = \frac{3600}{2 \times 60} \text{ Hz} \Rightarrow$

$$\lambda = \frac{v}{n} = \frac{760}{30} = 25.3 \text{ m.}$$

### [CHEMISTRY]

21. B  
22.

Reactivity of alkali metals:  $\text{Li} < \text{Na} < \text{K} < \text{Rb} < \text{Cs}$ .

Reactivity of halogens:  $\text{Fe} > \text{Cl} > \text{Br} > \text{I}$

- 23.

It is as per their emf values. The reducing power does not only depend upon ionisation energy but also on enthalpy of atomisation and enthalpy of hydration also.


24. A  
25.

The maximum covalency of Be is 4, e.g.,  $\text{Na}_2[\text{Be}(\text{OH})_4]$  while that of Al is 6, e.g.,  $\text{Na}_3[\text{AlF}_6]$ .


26. A  
27. B  
28.

IE of Mg:  $\text{Na} < \text{Al} < \text{Mg}$


- 29.




- 30.



31. A  
32.



33. D  
34.



35. Cs because of its low IE emits electron under the influence of even candle light.
36.  $\text{LiHCO}_3$  is unstable and exists only in solution.
37. Non-metal oxides being acidic decompose carbonates to evolve  $\text{CO}_2$  gas.
38.  $\text{NaNO}_3$  is called chile salt petre.
39. Mixture of  $\text{K}_2\text{CO}_3$  and  $\text{Na}_2\text{CO}_3$  is called fusion mixture.
40.  $\text{BaCO}_3 > \text{SrCO}_3 > \text{CaCO}_3 > \text{MgCO}_3$ . Thermal stability decreases as the basic character of the metal hydroxide decreases.